

Erratum: CP asymmetry in heavy Majorana neutrino decays at finite temperature: the nearly degenerate case

S. Biondini,^a N. Brambilla,^{a,b} M.A. Escobedo^c and A. Vairo^a

^a*Physik-Department, Technische Universität München,
James-Frank-Str. 1, 85748 Garching, Germany*

^b*Institute for Advanced Study, Technische Universität München,
Lichtenbergstrasse 2 a, 85748 Garching, Germany*

^c*Institut de Physique Théorique, Université Paris Saclay, CNRS, CEA,
F-91191 Gif-sur-Yvette, France*

E-mail: simone.biondini@tum.de, nora.brambilla@tum.de,
miguel-angel.escobedo-espinoza@cea.fr, antonio.vairo@tum.de

ERRATUM TO: JHEP03(2016)191

ARXIV EPRINT: [1511.02803](https://arxiv.org/abs/1511.02803)

We correct an error in the SU(2) part of (B.25), (B.26), (B.27). The equations should read

$$\text{Im}(-i\mathcal{D}_{3,\text{fig.23}}^\ell) + \text{Im}(-i\mathcal{D}_{4,\text{fig.23}}^\ell) = -\frac{\text{Im}[(F_1^* F_2)^2]}{(16\pi)^2 M} \frac{g'^2}{4} \left(1 - \frac{\Delta}{M}\right) \delta^{\mu\nu} \delta_{mn} + \dots, \quad (\text{B.25})$$

$$\text{Im}(-i\mathcal{D}_{1,\text{fig.24}}^\ell) + \text{Im}(-i\mathcal{D}_{2,\text{fig.24}}^\ell) = -\frac{\text{Im}[(F_1^* F_2)^2]}{(16\pi)^2 M} \frac{g'^2}{4} \left(-1 + \frac{\Delta}{M}\right) \delta^{\mu\nu} \delta_{mn} + \dots, \quad (\text{B.26})$$

$$\text{Im}(-i\mathcal{D}_{3,\text{fig.24}}^\ell) + \text{Im}(-i\mathcal{D}_{4,\text{fig.24}}^\ell) = -\frac{\text{Im}[(F_1^* F_2)^2]}{(16\pi)^2 M} \frac{g'^2}{4} \left(1 - \frac{\Delta}{M}\right) \delta^{\mu\nu} \delta_{mn} + \dots \quad (\text{B.27})$$

Note that the contributions from the SU(2) gauge bosons vanish for the diagrams of figure 23 and 24. The error influences our final results in the parts depending on the SU(2) gauge coupling g . Equations (4.1) and (4.2) for the matching coefficients should read

$$\begin{aligned} \text{Im} a_{11}^\ell = -\text{Im} a_{11}^{\bar{\ell}} = & \frac{\text{Im}[(F_1^* F_2)^2]}{(16\pi)^2} \left\{ 6\lambda \left[1 + \ln 2 - (2 - \ln 2) \frac{\Delta}{M} \right] \right. \\ & \left. - \frac{3}{8} g^2 \left[2 - \ln 2 + (3 - 5 \ln 2) \frac{\Delta}{M} \right] - \frac{g'^2}{8} \left[4 - \ln 2 + (1 - 5 \ln 2) \frac{\Delta}{M} \right] \right\}, \quad (4.1) \end{aligned}$$

$$\begin{aligned} \text{Im} a_{22}^\ell = -\text{Im} a_{22}^{\bar{\ell}} = & -\frac{\text{Im}[(F_1^* F_2)^2]}{(16\pi)^2} \left\{ 6\lambda \left[1 + \ln 2 + (2 - \ln 2) \frac{\Delta}{M} \right] \right. \\ & \left. - \frac{3}{8} g^2 \left[2 - \ln 2 - (3 - 5 \ln 2) \frac{\Delta}{M} \right] - \frac{g'^2}{8} \left[4 - \ln 2 - (1 - 5 \ln 2) \frac{\Delta}{M} \right] \right\}. \quad (4.2) \end{aligned}$$

As a consequence, equations (5.2), (5.3) and (5.6) for the unflavoured case should be

$$\Gamma_{11,\text{direct}}^{\ell,T} - \Gamma_{11,\text{direct}}^{\bar{\ell},T} = \frac{\text{Im}[(F_1^* F_2)^2]}{64\pi^2} \left\{ \lambda \left[1 + \ln 2 - (2 - \ln 2) \frac{\Delta}{M} \right] - \frac{g^2}{16} \left[2 - \ln 2 + (3 - 5 \ln 2) \frac{\Delta}{M} \right] - \frac{g'^2}{48} \left[4 - \ln 2 + (1 - 5 \ln 2) \frac{\Delta}{M} \right] \right\} \frac{T^2}{M}, \quad (5.2)$$

$$\epsilon_{1,\text{direct}}^T = \frac{\text{Im}[(F_1^* F_2)^2]}{8\pi|F_1|^2} \left(\frac{T}{M} \right)^2 \left\{ \lambda \left[2 - \ln 2 + (1 - 3 \ln 2) \frac{\Delta}{M} \right] - \frac{g^2}{16} \left[2 - \ln 2 + (3 - 5 \ln 2) \frac{\Delta}{M} \right] - \frac{g'^2}{48} \left[4 - \ln 2 + (1 - 5 \ln 2) \frac{\Delta}{M} \right] \right\}, \quad (5.3)$$

$$\epsilon_{2,\text{direct}}^T = - \frac{\text{Im}[(F_1^* F_2)^2]}{8\pi|F_2|^2} \left(\frac{T}{M} \right)^2 \left\{ \lambda \left[2 - \ln 2 - (9 - 5 \ln 2) \frac{\Delta}{M} \right] - \frac{g^2}{16} \left[2 - \ln 2 - 7(1 - \ln 2) \frac{\Delta}{M} \right] - \frac{g'^2}{48} \left[4 - \ln 2 - (9 - 7 \ln 2) \frac{\Delta}{M} \right] \right\}, \quad (5.6)$$

and similarly equations (7.9) and (7.10) for the flavoured case should read

$$\epsilon_{f1,\text{direct}}^T = \frac{\text{Im}[(F_1^* F_2)(F_{f1}^* F_{f2})]}{8\pi|F_1|^2} \left(\frac{T}{M} \right)^2 \left\{ \lambda \left[2 - \ln 2 + (1 - 3 \ln 2) \frac{\Delta}{M} \right] - \frac{g^2}{16} \left[2 - \ln 2 + (3 - 5 \ln 2) \frac{\Delta}{M} \right] - \frac{g'^2}{48} \left[4 - \ln 2 + (1 - 5 \ln 2) \frac{\Delta}{M} \right] \right\}, \quad (7.9)$$

$$\begin{aligned} \epsilon_{f2,\text{direct}}^T = & - \frac{\text{Im}[(F_1^* F_2)(F_{f1}^* F_{f2})]}{8\pi|F_2|^2} \left(\frac{T}{M} \right)^2 \left\{ \lambda \left[2 - \ln 2 - (9 - 5 \ln 2) \frac{\Delta}{M} \right] - \frac{g^2}{16} \left[2 - \ln 2 - 7(1 - \ln 2) \frac{\Delta}{M} \right] - \frac{g'^2}{48} \left[4 - \ln 2 - (9 - 7 \ln 2) \frac{\Delta}{M} \right] \right\} \\ & + \frac{\text{Im}[(F_1 F_2^*)(F_{f1}^* F_{f2})]}{2\pi|F_2|^2} \left(\frac{T}{M} \right)^2 \lambda \frac{\Delta}{M}. \end{aligned} \quad (7.10)$$

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.